



A comprehensive open-source course for teaching applied hydrological modelling in Central Asia

Beatrice Sabine Marti¹, Aidar Zhumabaev¹, and Tobias Siegfried¹

¹hydrosolutions GmbH, Zurich, Switzerland

Correspondence: Beatrice Marti (marti@hydrosolutions.ch)

Abstract. Numerical modelling is often used for climate impact studies in water resources management. It is, however, not yet highly accessible to many students of hydrology in Central Asia. One big hurdle for new learners is the scripting requirement for the preparation of relevant data prior to the actual modelling. We present a robust, open-source workflow and comprehensive teaching material that aims at bridging the gap between theoretical hydrological modelling know how and applied modelling for climate impact studies in Central Asia. The teaching material has been refined during two consecutive years and is being taken up by several professors teaching hydrological modelling in Central Asia.

1 Introduction

As sustainable water resources management will be ever more demanding in a changing world, the general need to strengthen the education of future water resources specialist does not subside (Vincent-Lancrin, 2009). Teaching future leaders in water resources management about hydrological modelling strengthens their capacity to find answers to the present and future challenges in using hydrological models effectively for decision making (Seibert and Vis, 2012; Wagener et al., 2012).

The Central Asia water sector is especially vulnerable to suffer from climate change impacts and there is not yet sufficient local capacity to adequately deal with the present and upcoming challenges (Abdullaev and Rakhmatullaev, 2016b). Much of the hydrological modelling work is currently done by foreign experts with sometimes questionable quality and limited transfer to local experts (Abdullaev and Rakhmatullaev, 2016a). Traditionally, Central Asian National Meteorological and Hydrological Services (CA NMHS) (organizations responsible for seasonal forecasts of water availability, among other tasks) rely on time series of station data for their predictions (Unger-Shayesteh et al., 2013) but the use of numerical models to support forecasts is increasingly researched (Gerlitz et al., 2020). Further, the collection and processing of hydrology-relevant data is the basis for robust modelling and capacities are being built up in Central Asia (Abdullaev and Rakhmatullaev, 2014). Thereby, the focus is not only on station data but also on remote sensing and re-analysis products. Modelling efforts might thus support the closing of knowledge gaps in linking observed runoff changes to changes in climatic and cryospheric changes in many headwater basins in Central Asia (Unger-Shayesteh et al., 2013).

Considerable efforts are put into the preparation of teaching resources by hydrologists worldwide and sharing platforms have been established to facilitate teaching. Examples are HydroShare (CUAHSI, 2022) and MOCHA (Wagener et al., 2012). Also worth mentioning is the Open Global Glacier Model Edu site which provides ample resources for teaching glacier dynamics



modelling (OGGM Consortium, 2022) or airGR (Coron et al., 2017), an open-source conceptual hydrological model which comes with elaborate online teaching functionality. In recent years, numerical modelling exercises are increasingly shared in the teaching community (e.g. Knoben and Spieler (2022), Seibert and Vis (2012)). However, the focus lies on general hydrology or US hydrology and comprehensive material on Central Asian hydrology is not readily available.

30 The aim of this paper is (1) to present a reproducible workflow for hydrological modelling for climate impact studies in Central Asia and (2) to provide open-source materials for teaching this workflow to students of hydrology and water resources management. The workflow and materials have been developed for a 2-week block course on applied hydrological modelling at the German-Kazakh University (DKU) in Almaty.

2 Teaching goals

35 Course participants understand the most important hydrological processes at the regional scale that are relevant in Central Asia and how future changes in climate and land use may impact their societies. They understand the differences between empirical, conceptual and physical hydrological models, the data requirements and properties of these model types as well as advantages and disadvantages of these model types. They know how such models are applied in the region by the CA NMHS for the reduction of risks from hydrological extremes and for improved water resources management for reservoir operations and irrigation scheduling. Course participants further know where to find high quality, publicly available data and can use open-source software effectively for catchment characterization. They can use software to implement a hydrological model of a catchment and they apply state-of-the art procedures to calibrate and validate a hydrological model. Last but not least, course participants can use hydrological models to simulate the impact of climate change on the discharge of a basin and present it to a scientific community.

45 3 Prerequisites

The master level course requires basic understanding of mathematics and computer skills. Prior knowledge of hydrology, R and QGIS are an advantage. Participation in the course requires a personal computer not older than about 5 years running Windows 10 or 11. All data to reproduce the examples in the on-line course book is available online (see section data and code availability). However, the bulk download requires approximately 36 GB of free storage space. If storage space is an issue, it is recommended to download only the data required for selected examples or exercises.

4 Software used

To ensure free access to the course material for all interested people, open-source/open-access software has been used in this course where possible. The required software includes the statistical programming language R version 4.2.1 (R Core Team, 2022) which is accessed through the graphical user interface RStudio (version 2022.07.1) (RStudio Team, 2022) and QGIS 3.24 (QGIS Development Team, 2022) with utilities of SAGA (Conrad et al., 2015) and GRASS (GRASS Development



Team, 2022). If suitable software was not available open-source, free, publicly available software was chosen, i.e. the hydraulic hydrological modelling software RS MINERVE (version 2.9.1.0) (Roquier et al., 2021). RS MINERVE implements the models HBV (Bergström, 1976) and GSM (Schaeffli et al., 2005), among others. RS Minerve runs on Windows OS only.

The course book is written using the Quarto authoring system (Quarto Development Team, 2022) and includes numerous code examples that facilitate the up-take of scripting skills for hydrological modelling. It is hosted publicly on GitHub and rendered via GitHub pages (https://github.com/hydrosolutions/caham_book). Readers can post issues directly on the GitHub Issues tracker. Apart from facilitated editing, the browser-based rendering has the advantage of automatic translation options available in many internet browsers nowadays.

The R package `riversCentralAsia` (Marti and Siegfried, 2022) was developed to support data processing tasks.

Moodle (Moodle project, 2022) was available from DKU for course administration and for depositing course material and video recordings of the lectures for the students. Lectures took place on-line via Zoom (Zoom Video Communications, Inc., 2022).

5 Data

All data used in the course can either be downloaded through public links in the course book or from public sources. Discharge monitoring locations and discharge time series at decadal (10-day) or monthly intervals for Kazakh, Kyrgyz, Tajik and Uzbek river basins are available from Hydrological Yearbook data (Uzbekhydromet, 2022).

Glacier outlines from the Randolph Glacier Inventory v6.0 (RGI) (RGI Consortium, 2017) were used to estimate glaciated areas of basins and glacier volumes. Glacier thinning rates (Hugonnet et al., 2021) and glacier ablation rates Miles et al. (2021) were used to manually calibrate the GSM model for glacier discharge. Estimates of daily snow water equivalent (SWE) between 2000 and 2011 from the High Mountain Snow Reanalysis product (Liu et al., 2021b) were used for model calibration.

The NASA SRTM digital elevation model 1 Arc-second (30 m) global product is used as a DEM (NASA JPL, 2013). Land cover information can be obtained from the Copernicus Global Land Service: Land Cover 100m: collection 3: epoch 2019: Globe data (Buchhorn et al., 2020). For basin characterization we use the CHELSA v2.1 climatologies (Karger et al., 2017) available for the periods 1981 - 2010, 2011 - 2040, 2041 - 2070 and 2071 - 2100. As the monitoring network in the Central Asia high mountain region is sparse (Unger-Shayesteh et al., 2013), bias-corrected re-analysis data is often used to drive hydrological models (e.g. Siegfried et al. (2012), Didovets et al. (2021), Khanal et al. (2021)). The course uses the CHELSA v2.1 data set (Karger et al., 2017, 2021b) at high resolution which contains daily precipitation and temperature time series at 1 km grid resolution globally between January 1979 and December 2011. CHELSA v2.1 time series are derived from the ERA5 re-analysis collection but perform better in mountainous regions because they are corrected for elevation biases (Beck et al., 2020). Future climate projections are downloaded for 4 shared socio-economic development pathways (SSP) (Riahi et al., 2017a), namely SSP1, SSP2, SSP3 and SSP5, for the 4 global circulation models (GCM) with the highest priorities according to the ISIMIP3b protocol (ISIMIP3 consortium, 2022) in the Coupled Model Inter-comparison Project (CMIP) phase 6, namely GFDL-ESM4 (Krasting et al., 2018), IPSL-CM6A-LR (Boucher et al., 2018), MRI-ESM2.0 (Yukimoto et al.,




Figure 1. Model chain for CMIP 6 climate impact analysis.

2019) and UKESM1.0-LL (Tang et al., 2019). The GCM model results are downscaled to the CHELSA v2.1 data using quantile mapping (Gudmundsson, 2016). 4 climate change scenarios, represented by 4 GCM models each are thus available for model forcing.

6 Model workflow

In a fast-evolving world, novel data sets describing weather or snow conditions for example, or more powerful software solutions become available at ever higher quality and frequency, requiring an adaptation of previously established workflows. The present modelling workflow for climate impact studies in Central Asian headwater basins (see Fig. 1) has been developed in 2021 and was updated in 2022.

6.1 Data pre-processing

One of the biggest hurdles to hydrological modelling is choosing and processing of the required data. This course therefore deliberately starts with the data preparation process and not directly with the hydrological modelling.

6.1.1 Discharge

The first step of the modelling process involves the review of the discharge data. Data gaps are filled with the long-term norm discharge of the respective time period. Basic discharge characteristics (average annual discharge, mean specific discharge) are computed. The R package riversCentralAsia has a number of functions to facilitate discharge characterization. For example the aggregation of time series data to warm and cold seasons and to hydrological years or automated plots of deviations from norm discharge. Discharge data is formatted and printed in an RS MINERVE-readable format.



6.1.2 Geospatial data

RS MINERVE features a GIS environment which can be used for semi-automated model generation. This feature speeds up the model implementation for regional models with a large number of hydrological response units (HRU). This feature requires the preparation of consistent GIS layers for HRU, river segments and river junctions (Roquier et al., 2022). This section describes how geospatial data, namely digital elevation model (DEM) and gauge location, are processed to derive the GIS layers for semi-automated model generation.

The DEM (NASA JPL, 2013) is run through the GRASS GIS (GRASS Development Team, 2022) `r.fill.dir` sink-filling algorithm (Jenson and Domingue, 1988) implemented in QGIS (QGIS Development Team, 2022). Using the geo-location of the discharge gauge (obtained from Uzbekhydromet (2022) and manually validated) the catchment boundary is determined using the `r.watershed` and the `r.water.outlet` algorithms and the resulting raster layer of the catchment area is converted to a polygon (step-by-step instructions are available in the course book under the section ‘Geospatial data’, ‘Catchment delineation’).

Hydrological response units (HRUs) are often used in semi-distributed conceptual hydrological models (Pilz et al., 2017). HRUs represent zones of similar run-off formation processes (Flügel, 1995). Depending on which hydrological processes are dominant, HRUs are derived based on basin geology, land use, topography or a combination of the aforementioned. In mountainous areas in Central Asia, river discharge is dominated by snow melt and overland flow processes and a derivation of HRUs based on landscape topography, i.e. elevation was chosen. The derivation of elevation-based HRUs is done with the `Reclassify by Table` algorithm applied to the sink-filled DEM in QGIS. Elevation intervals of 200 m are chosen. The resulting raster is subsequently polygonized to obtain a vector layer. The fields of the resulting HRU layer have to be adapted to contain consistent entries for import to RS MINERVE.

If glaciers are present in the basin, the glacier area is intersected with the HRUs area because glacier discharge will be implemented as GSM model objects for each glaciated elevation band in RS MINERVE. To reduce the number of model objects, glacier volume and area are aggregated per elevation band in the HRU layer.

The SAGA GIS (Conrad et al., 2015) algorithm `Channel Network and Drainage Basins` implemented in QGIS is used to derive the GIS layers for river sections and junctions required for semi-automated model generation in RS MINERVE. The algorithm uses the parameter `River Network Level` to determine the resolution of the resulting river network. Iterative manual adjustment of the parameter leads to the desired resolution of the river network. Manual post-processing of the resulting river and junctions layers are necessary to remove springs (in the junctions layer) and to edit the attribute table for import to RS MINERVE.

For teaching in the short block course, basins are simplified to one junction (the outflow junction) to which all HRU connect. River routing is not covered. The course book, however, does cover the implementation of hydrological models with sub-basins in the examples. The derivation of the sub-basins layer is done using the SAGA algorithms `Flow Accumulation`, `Channel Network` and `Watershed Basins`. The `Channel Network Cutoff Value` is a parameter determining the level of detail in the sub-basin layer. Iterative adjustment of the parameters may be necessary to achieve satisfactory sub-basins



140 which are consistent with the junctions and river layer. The resulting sub-basins layer has to be intersected with the HRU layer to obtain HRUs for each sub-basin.

A QGIS model has been implemented which is compatible with QGIS version 3.24 and which is available in the student case study pack. The derivation of HRUs can, however, also be done using the function `gen_basinElevationBands` of the `riversCentralAsia` package (Marti and Siegfried, 2022).

6.1.3 Climate data

145 Daily average temperature and precipitation for each HRU are extracted from the CHELSA v2.1 time series (Karger et al., 2021a) and reformatted for import to an RS MINERVE database using the function `gen_HRU_Climate_CSV_RSMinerve` of the R package `riversCentralAsia`. The CHELSA v2.1 data set between January 1, 1979 and December 31, 2011 serves as observation data.

150 Projected climate data of SSP1, SSP2, SSP3 and SSP5 (Riahi et al., 2017b) and 4 GCM projections for temperature and precipitation Krasting et al. (2018); Boucher et al. (2018); Yukimoto et al. (2019); Tang et al. (2019) are pre-processed and also extracted for each HRU. Climate projections are downscaled to the basin using quantile mapping (Gudmundsson, 2016) whereby the CHELSA v2.1 time series serve as reference observation data. Also, climate projections are written to RS MINERVE readable files.

6.2 Hydrological modelling

155 Having covered all the data pre-processing steps, the semi-automatic model generation in RS MINERVE is used to implement the hydrological model. Detailed step-by-step instructions are given in the course book (Siegfried and Marti, 2022b). The non-glaciated HRUs are assigned HBV (Bergström, 1976) model objects and the glaciated HRUs are assigned GSM Schaeffli et al. (2005) model objects. Evaporation is calculated by RS MINERVE using the Turc method (Roquier et al., 2022).

160 Because the historical observation period between 1979 and 2011 is relatively short, initial conditions are estimated using a model spin-up of the observation period. For subsequent simulations, a spin-up period of 1 year is implemented to reduce the impact of possibly sub-optimal initial conditions. The calibration period is set to the years 2000 to 2011 to take advantage of the availability of glacier elevation change (Hugonnet et al., 2021), glacier ablation (Miles et al., 2021) and SWE data from the HMASR data set (Liu et al., 2021a). The validation period is set to 1980 to 2000.

165 The calibration process is started with one parameter set for the entire model. That means that we do not initially allow for different parameter values in the HRU. If acceptable model performance cannot be achieved in this way, a zonation is applied, allowing for example for larger groundwater storage in the HRUs in the valley bottoms or for lower snow melt rates at very high elevations. Care is taken to avoid over-parameterization. RS MINERVE allows a very flexible composition of the objective function: Multiple model performance indicators are available which can be linearly combined with user-selected weights (Roquier et al., 2022). In the suggested workflow, the calibration process is started with an equal-weight of the general
170 performance indices Nash efficiency and Root-Mean-Squared-Error. Should in subsequent calibration cycles become apparent,



that for example minimum flows are not well represented, the optimization function is adapted to better suited indices like for example in the case of low flows the log-Nash efficiency.

The parameters GSM model objects are calibrated manually by comparing simulated changes in glacier thickness to observed changes in glacier thickness by Hugonnet et al. (2021), aggregated per elevation band. We further compare simulated glacier discharge rates with RS MINERVE to simulated glacier ablation rates provided by Miles et al. (2021). As the GSM model implemented in RS MINERVE assumes an infinite glacier reservoir to melt from, the modeller has to manually set glacier melt to zero when the glacier volume becomes zero, i.e. when the glacier has melted. For simple, linear models, this can be done as a post-processing step. In more complex models, the hydrological model has to be stopped at regular time intervals to turn off glacier melt if necessary.

The parameters of the snow modules of the HBV model objects are calibrated using the HMASR SWE product (Liu et al., 2021a). As RS MINERVE currently does not support automated calibration for SWE, manual calibration is done.

Calibration subsequently proceeds from the soil humidity reservoir to the upper soil reservoir and finally to the lower soil reservoir. Only discharge at the outlet point is available to constrain parameters of these 3 HBV reservoirs in all HRUs.

The validation period is used to verify the ability of the hydrological model to simulate discharge produced with hitherto unseen forcing. Resulting parameters are compared to the literature.

6.3 Scenario analysis

The calibrated model is now used to perform simulations of river discharge under the 16 climate change scenarios (4 GCM, 4 SSP). Simulation results are aggregated to 20-year average hydrographs of today (2000-2020), the middle of the century (2040-2060) and the end of the century (2080-2100) for each SSP.

190 7 Teaching modules

The learning goals are structured into theoretical components where students learn the basic concepts required for modelling and applied components where students learn hands-on how to implement a hydrological model. Students thereby follow the above described modelling workflow which is described in higher detail also in the course book (Siegfried and Marti, 2022b).

7.1 Modus operandi

Due to travel restrictions, the course was planned as an intensive 8-day on-line block course for master students of integrated water resources management at the German-Kazakh University (DKU) in Almaty. The course was taught remotely in spring 2021 and 2022 as an elective module at DKU for groups of approximately 12 students each. Two training of trainer courses were conducted in Spring 2022 for course transfer. 25 academic staff from 19 Central Asia higher education institutions attended the trainings.

The course will continue to be taught at DKU by a local scientific expert who participated in the students course twice. The course schedule consists of 3 lessons of 1 hour each of daily lectures, exercise discussions and case study presentations.



	Day 1	Day 2	Day 3	Day 4
Learning Question	<i>Characteristics of the Central Asian Region & role of hydrological modeling</i>	<i>What is a hydrological model and what are key hydrological processes?</i>	<i>What are data requirements for hydrological modeling?</i>	<i>What are data requirements for hydrological modeling?</i>
	Welcome & Housekeeping	Recap & homework discussion	Recap & homework discussion	Recap & homework discussion
1h	LECTURE Course prerequisites, discussion of assignments and student case studies	LECTURE Types of hydrological models (empirical, conceptual and physical models)	LECTURE - Water Balance	LECTURE Input data requirements for hydrological modeling (Discharge Data)
20min	Break	Break	Break	Break
1h	LECTURE Physical Geography, irrigation & hydropower, international relations	LECTURE Hydrological Processes	LECTURE Input data requirements for hydrological modeling (GIS Data)	LECTURE Input data requirements for hydrological modeling (Climate Data)
20min	Break	Break	Break	Break
1h	LECTURE Role of hydrological models to help address water-related challenges	LECTURE Hydrological Processes	LECTURE Input data requirements for hydrological modeling (GIS Data)	LECTURE - Modeling Process - Perceptual Model
Homework	PREP. WORK: Installation of software, download of case study packs	Homework Role play Model Application	EXERCISE 1: GIS Catchment Characterisation	EXERCISE 2: Characterisation of discharge time series
	Day 5	Day 6	Day 7	Day 8
Learning Question	<i>How does a hydrological model work?</i>	<i>How to find a good set of model parameters?</i>	<i>How to write a conference abstract?</i>	<i>How to present in front of an audience?</i>
	Recap & homework discussion	Recap & homework discussion	Recap & homework discussion	Final Exam Day
1h	LECTURE - Linear Reservoir Model - HBV Hydrological Model	LECTURE Model calibration and validation	SEMINAR Online group work with Q&A on exercises and final exam	Final Exam [All] Student Conference with presentation of case studies
20min	Break	Break	Break	Break
1h	LECTURE - Intro to RSMInerve	LECTURE Model calibration and validation	SEMINAR Online group work with Q&A on exercises and final exam	Final Exam [All] Student Conference with presentation of case studies
20min	Break	Break	Break	Break
1h	SEMINAR - Intro to RSMInerve - Presentation of Badam Model	DEMONSTRATION Model Calibration and validation	SEMINAR Online group work with Q&A on exercises and final exam	Course Wrap-Up [All] Course review questionnaire and oral course feedback by students
Homework	Homework RSMInerve User Manual: Example 1	EXERCISE 3: RSMInerve model implementation	EXERCISE 4: Submission of Conference Abstract	

Figure 2. A proposed course schedule for a 8 day course.

The students have been asked to do daily homework, including 4 graded exercises which, together with the individual course participation and a final exam, form the final grade for the course. Figure 2 shows the course schedule for the 8-day block teaching. In the first teaching round, a seminar with representatives from the private industry and the Central Asian Hydromets were invited to present the use hydrological modelling in their work. This seminar was very well received. A regional holiday did unfortunately not allow for such a seminar in the second course and it is not visible in the schedule in Fig. 2. The individual course modules are described in more detail in the following sections.

7.2 Data management

The data required for students to achieve the course requirements is made available online in a Dropbox repository (Siegfried and Marti, 2022a). Students have access to data to reproduce the examples provided in the course book and for modelling a case study basin. The below section describes how the data is organized.




Figure 3. Overview over the student case study basins. Background image: Stamen Terrain Background via QGIS quick map services. Map tiles by Stamen Design, under CC BY 3.0 using data by OpenStreetMap (© OpenStreetMap contributors 2022. Distributed under the Open Data Commons Open Database License (ODbL) v1.0.). River network and surface water bodies from Global Runoff Data Centre, GRDC (2020).

7.2.1 Student case study packs

Students work in groups of two on one of 8 case study basins in Central Asia (Chon Kemin, Ala Archa, Nauvalisoy, Pskem, Chatkal, Isfayramsoy, Sokh and Gunt from north to south in Fig. 3) for which data and code snippets have been prepared.

215 The data available for students includes CHELSA v2.1 climatologies for Central Asia (temperature T (Karger et al., 2017) and precipitation P (Beck et al., 2020)) as well as potential evaporation E_{pot} (Trabucco and Zomer, 2019) and the aridity index $\phi = E_{pot}/P$. Students further receive a code folder with scripts for the preparation of climate files and for the discharge characterization. In the folder ‘GaugeData’ students find the measured discharge in the tabular format as comma-separated values. These files can be read to R using the R package `riversCentralAsia` (Marti and Siegfried, 2022). The ‘GIS’ folder
220 includes a DEM which is already re-sampled to 100 m grid resolution and which was run through a sink-filling algorithm (Wang and Liu, 2006). The folder further contains a shapefile with point information about the gauge station at the outlet of the catchment and a shapefile with the boundaries of the catchment. To make sure also students with no prior experience in R scripting achieve the goal of setting up a hydrological model, necessary input layers for automatic model generation are made available as well. These include a layer with hydrological response units, a junctions and a river layer.

225 Glacier outlines (RGI Consortium, 2017) are provided in the ‘Glaciers’ folder. Further there is land cover information available (Buchhorn et al., 2020) in the ‘Landuse’ folder. Because the download and processing of the climate input data requires fast internet, considerable free storage space and computing power, climate inputs were pre-produced for the students



and are available in the folder RS_MINERVE. The data includes a file labeled hist_obs, referring to the historical observation period, 4 files hist_sim for each general circulation model (GCM) showing the GCM simulation results for the historical period
230 and 16 files fut_sim which contain forcing for 4 GCM and 4 shared socio-economic pathways.

7.2.2 Examples from the book

In order to reproduce examples from the course book, students are encouraged to download the required data from links provided in the course book. Data from the Gunt river in the Amu Darya Basin and Atbashy and Chirchiq rivers in the Syr Darya basin are available with the same naming conventions as in the student case study sets. Further data covering the entire
235 Central Asian region is available in the folder ‘central_asia_domain’ and contains a map of the irrigated area (Ragetti et al., 2018) under agriculture, historical (Karger et al., 2017, 2021b) and future climatologies (Krasting et al., 2018; Boucher et al., 2018; Yukimoto et al., 2019; Tang et al., 2019) in the ‘climatology’ folder, a land cover map (Buchhorn et al., 2020) in the ‘copernicus_land_cover’ folder, a geopackage layer of available geolocations of the most important discharge monitoring locations in Central Asia, obtained from Uzbekhydromet (2022) and manually validated, in the ‘gauges_basins’ folder and
240 glacier related data as glacier outlines of regions 13 (covering most of the river basins draining into Central Asia) of the Randolph Glacier Inventory (RGI) version 6 (RGI Consortium, 2017), glacier thickness by Farinotti et al. (2019) for RGI region 13, per-glacier elevation change rates by Hugonnet et al. (2021) and glacier balance and imbalance ablation rates by Miles et al. (2021) in the ‘glaciers’ folder.

7.3 Housekeeping

245 The course starts with an introduction of lecturers and course participants whereby the participants state their background and their previous experiences with hydrological modelling, R or QGIS. The main tutor then explains the course schedule and modus operandi during lectures. Further, the learning resources and software are introduced. The course material includes data, an R package with functions supporting the processing of the data, a course book, lecture slides and video tutorials. As the student groups were small, a highly interactive teaching style could be applied where course participants were encouraged
250 to actively participate in the lectures. The students form groups of 2 and select a case study basin on which they will work during the block course. Data and code snippets have been prepared by the teachers for each case study basin to make sure that students have the necessary data for modelling available, even if they should fail a preparatory exercise.

Every day starts with a recap session where students summarize what they have learned the previous day. Further, the homework and exercises are discussed and general questions are answered.

255 7.4 Day 1: Hydrological characteristics of the Central Asian region and role of hydrological modeling

After the introduction to the course, participants get a lecture on the physical geography of Central Asia, the irrigation and hydropower sectors and implications of water resources on international relations. In a second lecture block, participants learn about the role of hydrological models in helping to address water challenges.



As homework, course participants are asked to install the required software and to familiarize themselves with the software.

260 **7.5 Day 2: What is a hydrological model and what are are key hydrological processes?**

After the recap on day 2, course participants get an overview over the types of hydrological models (empirical, conceptual and physical) and how they can be used to address water challenges. In the second and third course block, students learn about the most important hydrological processes, watershed delineation, water balance as well as an overview over the dominant processes in the cryosphere.

265 As homework, course participants are asked to slip into the role of water professionals (3 profiles are provided). They describe in their own words the water challenges they are tasked to address and what information they require from a hydrological model. They then determine the type of hydrological model and model input data they require to inform their work. The homework is discussed in plenum the following day. Through this thinking exercise, students deepen their understanding about hydrological models and their applications.

270 **7.6 Days 3 and 4: What are data requirements for hydrological modeling?**

The third module dives into the data preparation part of the model chain (see Fig. 1). This part covers the processing of geospatial data (GIS layers), including the delineation of a basin and the derivation of the river network based on the DEM that can be used for automated model generation in RS MINERVE.

As homework, course participants work on their first graded exercise: The basin characterization using GIS data. Using data
275 provided in the student case study sets students extract basin characteristics and interpret their results in a short report. Step-by-step instruction videos, embeded in the on-line course book, are available that support first-time GIS users in this exercise. The videos are also available in a dedicated YouTube repository (Zhumabaev and Siegfried, 2022).

The module continues on day 4 of the course with lectures on discharge characterization, climate data and the modelling process. The presentation on discharge covers traditional hydrology like how to deal with data gaps and how to calculate basic
280 discharge statistics but also an outlook in the opportunities and difficulties introduced by automated discharge loggers. Students subsequently learn about data sources for climate forcing. The course participants learn how to extract time series of CHELSA climate data for all hydrological response units.

In the third lecture of this intensive course day, students are introduced to a general modelling process from conceptual model to model validation as described for example in Beven (2012). A key focus thereby is how model performance can be
285 judged. Thus, students are introduced to model performance indices. Students further learn about perceptual models of a basin following Wagener et al. (2007) and Gnann et al. (2021) and manually draw a perceptual model of their case study basin. This small in-class exercise builds on the basin characterization exercise (the previous homework) and helps the students to reflect on the hydrological processes they expect to be dominant in their basins.

As homework, course participants analyse the discharge time series of their chosen catchment and write up a short report
290 for their second graded exercise. Students can copy code snippets from the course book or use a prepared R script available in



the student case study pack for producing a selection of figures to characterize discharge. The interpretation of the figures has to be done by the students themselves.

7.7 Day 5: How does a hydrological model work?

Typically, the recap and discussion of the homework takes up a considerable amount of time on day 5. The lectures therefore are abbreviated. Students are introduced to the linear reservoir model and learn to differentiate between model states, variables and parameters. Then the HBV model (Bergström, 1976) as implemented in RS MINERVE (Roquier et al., 2022) is introduced. Subsequently, the software RS MINERVE is introduced in a online demonstration and a case study, the Badam reservoir system in South Kazakhstan, is presented. This case study demonstrates an advanced hydrological-hydraulic model of a natural river system that has been altered by different man-made reservoirs and inter-basin water transfers.

As homework, students go through the exercises in the RS MINERVE manual to familiarise themselves with the software.

7.8 Day 6: How to find a good set of model parameters?

Students learn about model calibration. Linear regression is introduced and model goodness of fit is discussed. Objective functions and global optimization are introduced as well as strategies to avoid over-parameterization. Further, students learn how to divide their available time series into warm-up, calibration, validation and baseline periods. The content is summarized in a calibration strategy.

As homework, students are asked to use the readily available GIS layers for automated model implementation in RS MINERVE. A tutorial video is available that demonstrates how to do this.

7.9 Day 7: Model calibration and how to write a scientific abstract?

Students have time to collaborate on-line and manually calibrate the hydrological model for their case study site. Tutors are also on-line and provide support where needed. Students further have time to finalize their presentations for the last day of the course. To support their work they receive presentation guide lines and a short demo presentation of a model of the Gunt river basin.

As third graded exercise, students have to hand in a calibrated model of their case study basin.

As homework, students write an abstract for the student conference which takes place on the last day of the course. They have to comply to abstract submission guide lines of the Silk Road of Knowledge conference. The abstract is the fourth graded exercise.

7.10 Day 8: Student conference and course wrap-up

Student groups take turns to present their hydrological modelling work of the past days in a pseudo-conference setting. Their presentations are graded. After the student conference, the students are asked to give feedback on the course face-to-face and in an anonymous Google form.



8 Experiences

The proposed workflow has been extensively tested and applied to further basins in Central Asia within the frame of Master thesis projects at ETH Zurich and DKU and also within the frame of the Hydro4U project (Hydro4U consortium, 2022). It provides a solid basis for hydrological modelling for climate change impact studies. As new data becomes available or new
325 model goals emerge (e.g. operational forecast), the above workflow needs to be adapted.

Students background at DKU was diverse with limited prior knowledge in scripting or numerical modelling. They found the course highly relevant to their field and interesting but also very challenging. Students rated the overall quality of the course exceptionally high with 4.9 out of 5 points. The worst grade (4.2 out of 5) was given to the homework tasks which were perceived as too difficult. This was reflected also in the rather poor quality of the homework handed in.

330 On-line teaching posed several challenges like poor internet connection of some students. Especially our Afghan students in the 2021 course suffered from frequent interruptions. Further, the installation of the software involved considerable off-hour tutoring need by some students. The use of university computers in the case where in-person teaching is possible or the use of RStudio cloud services could reduce the software installation hurdle.

The excellent reception of the modelling course led to two week-long training of trainers seminars for 25 academic staff from
335 19 different organizations throughout the region. The course was well received also by trainers and several plan to include at least part of the course material in their teaching program. As a spin-off of the training of trainers seminar, professors from the Kazakh National University, the Eurasian University in Kazakhstan, the Uzbek National University, and the German Kazakh University continue to work together over the summer by regularly meeting online for the discussion of relevant topics.

Authors as well as trainers agree that a block course is not sufficient for students to learn to master the tools required
340 for hydrological modelling. Separate courses on the use of QGIS and R should be taken before the start of the hydrological modelling course. Also, more time should be allowed for students to grasp the lecture content through further exercises. The course material should optimally be taught in a semester course.

9 Conclusions

Future hydrologists need to learn software and scripting tools as well as numerical hydrological modelling in addition to tradi-
345 tional hydrology as they are excessively used today in science but also in applied modelling. The presented course material and modelling workflow are open-source, readily available and tailored for climate impact studies in Central Asia. The robustness and efficacy of the workflow has been demonstrated in consulting tasks.

Code and data availability. The open-source course book is available via github https://hydrosolutions.github.io/caham_book/ (Siegfried and Marti, 2022b). The Link to the course book includes the link to the students case study packs which are available for download from
350 dropbox: <https://bit.ly/3CMWbbm>. All examples from the course book can be reproduced with the full CAHAM data package available from



here: <https://bit.ly/3e0tb5F> (attention, approx 36 GB size!). The R package extensively used in the course book is available for installation via github: <https://github.com/hydrosolutions/riversCentralAsia>. Lecture slides are available from dropbox: <https://bit.ly/3ReBVnx>.

Video supplement. Tutorial videos with step-by-step instructions are available from youtube: <https://bit.ly/3B2Emnr>

355 *Author contributions.* Tobias Siegfried designed the teaching course, prepared most of the teaching material and performed most of the teaching and exercise tutoring. Beatrice Marti contributed to the course exercises, sections on cryosphere data and hydrological modelling and took over said sections of the teaching. Aidar Zhumabaev recorded tutorial videos, supported students in exercises and presented his case study: the hydrological-hydraulic model of the Badam reservoir system.

Competing interests. The authors declare no competing interests.

360 *Acknowledgements.* The generous support by the Blue Peace Central Asia Programme at the Swiss Agency for Development and Cooperation is acknowledged. We especially thank Stéphanie Piers de Raveschoot, André Wehrli, Dimka Stantchev and Lisa Gampp there. At the German Kazakh University, Dr. Larissa Kogutenko is acknowledged for her amazing administrative and in-course support. We would further like to thank the authors of open-source software and data used in this article for making their work accessible.



References

- Abdullaev, I. and Rakhmatullaev, S.: Data Management for Integrated Water Resources Management in Central Asia, *Journal of Hydroinformatics*, 16, 1425–1440, <https://doi.org/10.2166/hydro.2014.097>, 2014.
- Abdullaev, I. and Rakhmatullaev, S.: River Basin Management in Central Asia: Evidence from Isfara Basin, Fergana Valley, *Environmental Earth Sciences*, 75, 677, <https://doi.org/10.1007/s12665-016-5270-9>, 2016a.
- Abdullaev, I. and Rakhmatullaev, S.: Setting up the Agenda for Water Reforms in Central Asia: Does the Nexus Approach Help?, *Environmental Earth Sciences*, 75, 870, <https://doi.org/10.1007/s12665-016-5409-8>, 2016b.
- 365 Beck, H. E., Wood, E. F., McVicar, T. R., Zambrano-Bigiarini, M., Alvarez-Garretón, C., Baez-Villanueva, O. M., Sheffield, J., and Karger, D. N.: Bias Correction of Global High-Resolution Precipitation Climatologies Using Streamflow Observations from 9372 Catchments, *Journal of Climate*, 33, 1299–1315, <https://doi.org/10.1175/JCLI-D-19-0332.1>, 2020.
- Bergström, S.: Development and Application of a Conceptual Runoff Model for Scandinavian Catchments, SMHI Report Nr Rho 7, Lund Institute of Technology/University of Lund, 1976.
- 375 Beven, K.: *Rainfall-Runoff Modelling*, John Wiley & Sons, Ltd, <https://doi.org/https://doi.org/10.1002/9781119951001>, 2012.
- Boucher, O., Denvil, S., Levvasseur, G., Cozic, A., Caubel, A., Foujols, M.-A., Meurdesoif, Y., Bony, S., Flavoni, S., Idelkadi, A., Mellul, L., Musat, I., and Saint-Lu, M.: IPSL IPSL-CM6A-LR Model Output Prepared for CMIP6 CFMIP, <https://doi.org/10.22033/ESGF/CMIP6.1522>, 2018.
- Buchhorn, M., Smets, B., Bertels, L., De Roo, B., Lesiv, M., Tsendbazar, N.-E., Herold, M., and Fritz, S.: Land Cover 100m: Collection 3: Epoch 2019: Globe 2020, DOI 10.5281/zenodo.3939050, 2020.
- 380 Conrad, O., Bechtel, B., Bock, M., Dietrich, H., Fischer, E., Gerlitz, L., Wehberg, J., Wichmann, V., and Böhner, J.: System for Automated Geoscientific Analyses (SAGA) v. 2.1.4, *Geoscientific Model Development*, 8, 1991–2007, <https://doi.org/10.5194/gmd-8-1991-2015>, 2015.
- Coron, L., Thirel, G., Delaigue, O., Perrin, C., and Andréassian, V.: The Suite of Lumped GR Hydrological Models in an R Package, *Environmental Modelling & Software*, 94, 166–171, <https://doi.org/10.1016/j.envsoft.2017.05.002>, 2017.
- CUAHSI: [hydroshare.org](https://www.hydroshare.org), <https://www.hydroshare.org/>, accessed: 2022-08-22, 2022.
- Didovets, I., Lobanova, A., Krysanova, V., Menz, C., Babagaliyeva, Z., Nurbatsina, A., Gavrilenko, N., Khamidov, V., Umirbekov, A., Qodirov, S., Muhyeyew, D., and Hattermann, F. F.: Central Asian Rivers under Climate Change: Impacts Assessment in Eight Representative Catchments, *Journal of Hydrology: Regional Studies*, 34, <https://doi.org/10.1016/j.ejrh.2021.100779>, 2021.
- 390 Farinotti, D., Huss, M., Fürst, J. J., Landmann, J., Machguth, H., Maussion, F., and Pandit, A.: A Consensus Estimate for the Ice Thickness Distribution of All Glaciers on Earth - Dataset, <https://doi.org/10.3929/ethz-b-000315707>, 2019.
- Flügel, W.-A.: Delineating hydrological response units by geographical information system analyses for regional hydrological modelling using PRMS/MMS in the drainage basin of the River Bröl, Germany, *Hydrological Processes*, 9, 423–436, <https://doi.org/10.1002/hyp.3360090313>, 1995.
- 395 Gerlitz, L., Vorogushyn, S., and Gafurov, A.: Climate Informed Seasonal Forecast of Water Availability in Central Asia: State-of-the-art and Decision Making Context, *Water Security*, 10, 100 061, <https://doi.org/10.1016/j.wasec.2020.100061>, 2020.
- Global Runoff Data Centre, GRDC: WMO Basins and Sub-Basins, https://www.bafg.de/GRDC/EN/02_srvcs/22_gslrs/gislayers_node.html, 2020.



- Gnann, S. J., McMillan, H. K., Woods, R. A., and Howden, N. J. K.: Including Regional Knowledge Improves Baseflow Signature Predictions in Large Sample Hydrology, *Water Resources Research*, 57, e2020WR028354, <https://doi.org/10.1029/2020WR028354>, 2021.
- GRASS Development Team: Geographic Resources Analysis Support System (GRASS GIS) Software, Version 8.2.1dev, Open Source Geospatial Foundation, <http://grass.osgeo.org>, 2022.
- Gudmundsson, L.: Qmap: Statistical Transformations for Post-Processing Climate Model Output, <https://CRAN.R-project.org/package=qmap>, 2016.
- 405 Hugonnet, R., McNabb, R., Berthier, E., Menounos, B., Nuth, C., Girod, L., Farinotti, D., Huss, M., Dussailant, I., Brun, F., and Käab, A.: Accelerated Global Glacier Mass Loss in the Early Twenty-First Century, *Nature*, 592, 726–731, <https://doi.org/10.1038/s41586-021-03436-z>, 2021.
- Hydro4U consortium: Hydro4U, <https://hydro4u.eu/>, accessed: 2022-09-01, 2022.
- ISIMIP3 consortium: ISIMIP3b simulation protocol, <https://protocol.isimip.org/>, accessed: 2022-08-30, 2022.
- 410 Jenson, S. K. and Domingue, J. O.: Extracting Topographic Structure from Digital Elevation Data for Geographic Information-System Analysis, *Photogrammetric Engineering and Remote Sensing*, 54, 1593–1600, 1988.
- Karger, D. N., Conrad, O., Böhner, J., Kawohl, T., Kreft, H., Soria-Auza, R. W., Zimmermann, N. E., Linder, H. P., and Kessler, M.: Climatologies at High Resolution for the Earth’s Land Surface Areas, *Scientific Data*, 4, 170122, <https://doi.org/10.1038/sdata.2017.122>, 2017.
- 415 Karger, D. N., Conrad, O., Böhner, J., Kawohl, T., Kreft, H., Soria-Auza, R. W., Zimmermann, N. E., Linder, H. P., and Kessler, M.: Climatologies at High Resolution for the Earth’s Land Surface Areas, *EnviDat*, <https://doi.org/10.16904/envidat.228.v2.1>, 2021a.
- Karger, D. N., Wilson, A. M., Mahony, C., Zimmermann, N. E., and Jetz, W.: Global Daily 1 Km Land Surface Precipitation Based on Cloud Cover-Informed Downscaling, *Scientific Data*, 8, 307, <https://doi.org/10.1038/s41597-021-01084-6>, 2021b.
- Khanal, S., Lutz, A., Kraaijenbrink, P. D. A., van den Hurk, B., Yao, T., and Immerzeel, W. W.: Variable 21st Century Climate Change Response for Rivers in High Mountain Asia at Seasonal to Decadal Time Scales, *Water Resources Research*, 57, e2020WR029266, <https://doi.org/10.1029/2020WR029266>, 2021.
- Knoben, W. J. M. and Spieler, D.: Teaching Hydrological Modelling: Illustrating Model Structure Uncertainty with a Ready-to-Use Computational Exercise, *Hydrology and Earth System Sciences*, 26, 3299–3314, <https://doi.org/10.5194/hess-26-3299-2022>, 2022.
- Krasting, J. P., John, J. G., Blanton, C., McHugh, C., Nikonov, S., Radhakrishnan, A., Rand, K., Zadeh, N. T., Balaji, V., Durachta, J., Dupuis, C., Menzel, R., Robinson, T., Underwood, S., Vahlenkamp, H., Dunne, K. A., Gauthier, P. P., Ginoux, P., Griffies, S. M., Hallberg, R., Harrison, M., Hurlin, W., Malyshev, S., Naik, V., Paulot, F., Paynter, D. J., Ploshay, J., Reichl, B. G., Schwarzkopf, D. M., Seman, C. J., Silvers, L., Wyman, B., Zeng, Y., Adcroft, A., Dunne, J. P., Dussin, R., Guo, H., He, J., Held, I. M., Horowitz, L. W., Lin, P., Milly, P., Shevliakova, E., Stock, C., Winton, M., Wittenberg, A. T., Xie, Y., and Zhao, M.: NOAA-GFDL GFDL-ESM4 Model Output Prepared for CMIP6 CMIP, <https://doi.org/10.22033/ESGF/CMIP6.1407>, 2018.
- 425 Liu, Y., Fang, Y., and Margulis, S. A.: High Mountain Asia UCLA Daily Snow Reanalysis, doi:<https://doi.org/10.5067/HNAUGJQXSCVU>, 2021a.
- Liu, Y., Fang, Y., and Margulis, S. A.: Spatiotemporal Distribution of Seasonal Snow Water Equivalent in High Mountain Asia from an 18-Year Landsat–MODIS Era Snow Reanalysis Dataset, *The Cryosphere*, 15, 5261–5280, <https://doi.org/10.5194/tc-15-5261-2021>, 2021b.
- Marti, B. and Siegfried, T.: riversCentralAsia: An R package for hydrological modelling, *Journal of Open-Source Software*, submitted, 2022.
- 435 Miles, E., McCarthy, M., Dehecq, A., Kneib, M., Fugger, S., and Pellicciotti, F.: Health and Sustainability of Glaciers in High Mountain Asia, *Nature Communications*, 12, 10, <https://doi.org/10.1038/s41467-021-23073-4>, 2021.



- Moodle project: Moodle 4.0, <https://moodle.org/>, accessed: 2022-08-24, 2022.
- NASA JPL: NASA Shuttle Radar Topography Mission Global 1 Arc Second [Data Set], Tech. rep., NASA EOSDIS Land Processes DAAC, 2013.
- 440 OGGM Consortium: Open Global Glacier Model Edu, <https://edu.oggm.org/en/latest/>, accessed: 2022-08-23, 2022.
- Pilz, T., Francke, T., and Bronstert, A.: lumpR 2.0.0: An R Package Facilitating Landscape Discretisation for Hillslope-Based Hydrological Models, *Geoscientific Model Development*, 10, 3001–3023, <https://doi.org/10.5194/gmd-10-3001-2017>, 2017.
- QGIS Development Team: QGIS Geographic Information System, <http://qgis.osgeo.org>, 2022.
- Quarto Development Team: Quarto, <https://quarto.org/>, accessed: 2022-09-01, 2022.
- 445 R Core Team: R: A Language and Environment for Statistical Computing, <https://www.R-project.org/>, 2022.
- Ragettli, S., Herberz, T., and Siegfried, T.: An Unsupervised Classification Algorithm for Multi-Temporal Irrigated Area Mapping in Central Asia, *Remote Sensing*, 10, 1823, <https://doi.org/10.3390/rs10111823>, 2018.
- RGI Consortium: Randolph Glacier Inventory – A Dataset of Global Glacier Outlines: Version 6.0, <https://doi.org/10.7265/N5-RGI-60>, 2017.
- 450 Riahi, K., van Vuuren, D. P., Kriegler, E., Edmonds, J., O’Neill, B. C., Fujimori, S., Bauer, N., Calvin, K., Dellink, R., Fricko, O., Lutz, W., Popp, A., Cuaresma, J. C., Kc, S., Leimbach, M., Jiang, L., Kram, T., Rao, S., Emmerling, J., Ebi, K., Hasegawa, T., Havlik, P., Humpenöder, F., Da Silva, L. A., Smith, S., Stehfest, E., Bosetti, V., Eom, J., Gernaat, D., Masui, T., Rogelj, J., Strefler, J., Drouet, L., Krey, V., Luderer, G., Harmsen, M., Takahashi, K., Baumstark, L., Doelman, J. C., Kainuma, M., Klimont, Z., Marangoni, G., Lotze-Campen, H., Obersteiner, M., Tabeau, A., and Tavoni, M.: The Shared Socioeconomic Pathways and Their Energy, Land Use, and Greenhouse Gas Emissions Implications: An Overview, *Global Environmental Change*, 42, 153–168, <https://doi.org/10.1016/j.gloenvcha.2016.05.009>,
- 455 2017a.
- Riahi, K., van Vuuren, D. P., Kriegler, E., Edmonds, J., O’Neill, B. C., Fujimori, S., Bauer, N., Calvin, K., Dellink, R., Fricko, O., Lutz, W., Popp, A., Cuaresma, J. C., Kc, S., Leimbach, M., Jiang, L., Kram, T., Rao, S., Emmerling, J., Ebi, K., Hasegawa, T., Havlik, P., Humpenöder, F., Da Silva, L. A., Smith, S., Stehfest, E., Bosetti, V., Eom, J., Gernaat, D., Masui, T., Rogelj, J., Strefler, J., Drouet, L., Krey, V., Luderer, G., Harmsen, M., Takahashi, K., Baumstark, L., Doelman, J. C., Kainuma, M., Klimont, Z., Marangoni, G., Lotze-Campen, H., Obersteiner, M., Tabeau, A., and Tavoni, M.: The Shared Socioeconomic Pathways and Their Energy, Land Use, and Greenhouse Gas Emissions Implications: An Overview, *Global Environmental Change*, 42, 153–168, <https://doi.org/10.1016/j.gloenvcha.2016.05.009>,
- 460 2017b.
- Roquier, B., Fluixa Sanmartin, J., Brauchli, T., Baracchini, T., Garcia Hernandez, J., Foehn, A., Paredes Arquiola, J., and De Cesare, G.: RS Minerve, <https://crealp.ch/rs-minerve/>, 2021.
- 465 Roquier, B., Fluixa Sanmartin, J., Brauchli, T., Baracchini, T., Garcia Hernandez, J., Foehn, A., Paredes Arquiola, J., and De Cesare, G.: RS Minerve Manuals, <https://crealp.github.io/rsminerve-releases/>, 2022.
- RStudio Team: RStudio: Integrated Development for R, <https://www.rstudio.com/>, 2022.
- Schaefli, B., Hingray, B., Niggli, M., and Musy, A.: A Conceptual Glacio-Hydrological Model for High Mountainous Catchments, *Hydrology and Earth System Sciences*, p. 15, 2005.
- 470 Seibert, J. and Vis, M. J. P.: Teaching Hydrological Modeling with a User-Friendly Catchment-Runoff-Model Software Package, *Hydrological Earth Syst. Sci.*, p. 11, <https://doi.org/10.5194/hess-16-3315-2012>, 2012.
- Siegfried, T. and Marti, B.: Data repository for the open-source course "Modeling of Hydrological Systems in Semi-Arid Central Asia", <https://bit.ly/3e0tb5F>, 2022a.



- 475 Siegfried, T. and Marti, B.: Hydrological Modelling in Semi-Arid Central Asia, hydrosolutions GmbH, Zurich, <https://doi.org/10.5281/zenodo.6349983>, 2022b.
- Siegfried, T., Bernauer, T., Guennet, R., Sellars, S., Robertson, A. W., Mankin, J., Bauer-Gottwein, P., and Yakovlev, A.: Will Climate Change Exacerbate Water Stress in Central Asia?, *Climate Change*, 112, 881–899, <https://doi.org/DOI.10.1007/s10584-011-0253-z>, 2012.
- Tang, Y., Rumbold, S., Ellis, R., Kelley, D., Mulcahy, J., Sellar, A., Walton, J., and Jones, C.: MOHC UKESM1.0-LL Model Output Prepared for CMIP6 CMIP Esm-piControl, <https://doi.org/10.22033/ESGF/CMIP6.5953>, 2019.
- 480 Trabucco, A. and Zomer, R.: Global Aridity Index and Potential Evapotranspiration (ET0) Climate Database V2, <https://doi.org/10.6084/m9.figshare.7504448.v3>, 2019.
- Unger-Shayesteh, K., Vorogushyn, S., Farinotti, D., Gafurov, A., Duethmann, D., Mandychew, A., and Merz, B.: What Do We Know about Past Changes in the Water Cycle of Central Asian Headwaters? A Review | Elsevier Enhanced Reader, *Global and Planetary Change*, 110, 4–25, <https://doi.org/10.1016/j.gloplacha.2013.02.004>, 2013.
- 485 Uzbekhydromet: Hydrological Yearbook data, 2022.
- Vincent-Lancrin, S.: The Role of Education in Capacity Development, in: *Capacity Development for Improved Water Management*, edited by Blokland, M. W., Alaerts, G., Kaspersma, J., and Hare, M., p. 24, CRC Press, London, first edn., 2009.
- Wagener, T., Sivapalan, M., Troch, P., and Woods, R.: Catchment Classification and Hydrologic Similarity, *Geography Compass*, 1, 901–931, <https://doi.org/10.1111/j.1749-8198.2007.00039.x>, 2007.
- 490 Wagener, T., Kelleher, C., Weiler, M., McGlynn, B., Gooseff, M., Marshall, L., Meixner, T., McGuire, K., Gregg, S., Sharma, P., and Zappe, S.: It Takes a Community to Raise a Hydrologist: The Modular Curriculum for Hydrologic Advancement (MOCHA), *Hydrology and Earth System Sciences*, 16, 3405–3418, <https://doi.org/10.5194/hess-16-3405-2012>, 2012.
- Wang, L. and Liu, H.: An Efficient Method for Identifying and Filling Surface Depressions in Digital Elevation Models for Hydrologic Analysis and Modelling, *International Journal of Geographical Information Science*, 20, 193–213, <https://doi.org/10.1080/13658810500433453>,
495 2006.
- Yukimoto, S., Koshiro, T., Kawai, H., Oshima, N., Yoshida, K., Urakawa, S., Tsujino, H., Deushi, M., Tanaka, T., Hosaka, M., Yoshimura, H., Shindo, E., Mizuta, R., Ishii, M., Obata, A., and Adachi, Y.: MRI MRI-ESM2.0 Model Output Prepared for CMIP6 CMIP Historical, <https://doi.org/10.22033/ESGF/CMIP6.6842>, 2019.
- Zhumabaev, A. and Siegfried, T.: Learning videos for the course: Modeling of Hydrological Systems in Semi-Arid Central Asia, <https://bit.ly/3B2Emnr>, 2022.
- 500 Zoom Video Communications, Inc.: Zoom, <https://zoom.us/>, accessed: 2022-08-25, 2022.