

# Brief communication: How deep is the snow at ~~the~~ Mount Everest?

Wei Yang<sup>1</sup>, Huabiao Zhao<sup>1</sup>, Baiqing Xu<sup>1</sup>, Jiule Li<sup>1</sup>, Weicai Wang<sup>1</sup>, Guangjian Wu<sup>1</sup>, Zhongyan Wang<sup>1</sup>, Tandong Yao<sup>1</sup>

<sup>1</sup>State Key Laboratory of Tibetan Plateau Earth System, Environment and Resources (TPESER), Institute of Tibetan Plateau Research, Chinese Academy of Sciences, Beijing 100101, China

Correspondence to: Huabiao Zhao (zhaohb@itpcas.ac.cn)

**Abstract.** Exploring the snow ~~depth~~thickness at ~~the~~ Mount Everest has long been a topic of interest ~~for studying geodesy, cryosphere and climate change in this inaccessible place on our planet. Previously reported snow depths have been inconsistent and have large uncertainties, but has not yet been measured successfully.~~ Here, we report the ground-penetrating radar survey of snow ~~depth~~thickness along the northern slope of ~~the~~ Mount Everest in May 2022. Our radar measurements display a gradual increasing transition of snow ~~depth~~thickness along the north slope, and the mean ~~snow depth~~snow thickness estimates at ~~the Mount the summit Everest is~~are approximately  $9.5 \pm 1.2$  m. This updated snow ~~depth~~thickness at ~~the~~ Mount Everest is much deeper than previously reported values (0.9~3.5 m).

## 1 Introduction

Mount Everest, ~~or~~ ~~Chomolungma,~~ ~~is one of~~ ~~the most inaccessible places on our planet~~the Earth summit, which is considered to be the most iconic peak (Kang et al., 2022; Matthews et al., 2020). There are very strong scientific and public motivations for determining the snow ~~depth~~thickness at ~~the~~ Mount Everest. Although China and Nepal jointly declared that the snow height of ~~the~~ Mount Everest ~~is was~~ 8848.86 metres above sea level (m asl) in 2020, the true rock height ~~has not been was not~~ precisely determined due to the unknown snow ~~depth~~thickness below. ~~And t~~The snow ~~depth~~thickness at extremely high elevations may vary dynamically with different seasons and ~~climate changes~~years (Potocki et al., 2022). Knowledge about snow depths during different periods will be helpful for explaining the discrepancies ~~of in the~~ reported snow heights at ~~the~~ Mount Everest, which ~~has have~~ been introduced by repeated surveys (Angus-Leppan, 1982; Chen et al., 2010; Xie et al., 2021). In additions, ~~snow layering at mountain summits contains information about local seasonal snow accumulation and climate history. However, snow and ice display accelerated loss rates in almost all regions on Earth~~ (Hugonnet et al., 2021; Kraaijenbrink et al., 2021). ~~S~~Similar to other snow/glacier-covered summits (Thompson et al., 2009), the snow and glaciers at Mount Everest are the sentinels for climate change and therefore offer a potential natural platform for understanding ongoing climate change at such extremely high elevations (Matthews et al., 2020; Potocki et al., 2022) and their possible widespread influence on the Asian Water Tower (Immerzeel et al., 2020). Comparisons of snow depth/stratigraphy during different periods may be potentially helpful for understanding the possible influence of anthropogenic climate change at extremely high elevations in the Himalayas (Brun et al., 2022; Pepin et al., 2022; Potocki et

al., 2022). Thus, the state of snow at the Mount Everest are critical for studies related to the geodesy, the cryosphere, and climate change.

35 However, previous attempts to measure the snow thickness at the Mount Everest were not successful due to harsh environment and instrument/method limitation. Previously reported snow depths derived by different methods and instruments ranged from 0.92 m to 3.5 m at Mount Everest. In 1975, a Chinese expedition team reported an estimated snow depth of 0.92 m by inserting a wooden stake into the snow (Chen et al., 2010). In 1992, a joint Chinese-Italian joint expedition team estimated a thickness of 2.52 m by inserting a steel stake into the snow (Chen et al., 2010). These results derived by stake methods were subjected to many factors such as snow density, stake length, and manpower issues at such harsh altitudes. Radio echo sounding is a suitable technique for imaging snow-ice environments and their internal structures (Rignot et al., 2013). In 2005, a Chinese mountaineering and surveying team claimed a snow depth of ~3.5 m by utilizing ground penetrating radar; however, the reported boundary between the snow and rock on the radar image was too ambiguous to provide an undisputed depth (Sun et al., 2006). In 2019 and 2020, various Nepalese and Chinese expedition teams measured the snow depth using different radar instruments; however, no results were reported (GSSI, 2021; Tone, 2020). Supported by the Second Tibetan Plateau Scientific Expedition and Research, we organized “The Earth Summit Missions 2022” expedition during the period from April to May 2022. One of our key goals is was to measure the snow depth at the Mount Everest.

## 2 Data and method

50 Ground penetrating radar (GPR) is a powerful tool in the field of cryosphere that has been widely used to survey snow depth (Holbrook et al., 2016; Yamamoto et al., 2004). To maximize portability at the Mount Everest, we conducted our GPR survey with a single transmitter–receiver antenna at a frequency of 1000 MHz using a Sensor & Software Pulse EKKO Pro system on May 4<sup>th</sup> 2022. In contrast with the previous radar survey conducted at the summit (Sun et al., 2006), our measurement started from the exposed metamorphosed limestone at an elevation that was approximately 15 m lower than the the summit–Mount Everest to ensure a gradual transition in the radar reflection profile and thus produce more easily post-55 discerning boundaries between the snow and rock (Fig. 1a). For all GPR measurementing points, a portable global navigation satellite system (UniStrong G138BD) continuously recorded the antenna locations. We obtained a total of 57 radar wavelet traces at irregularly spaced intervals (~0.5-1 m along the north slope and 0.2-0.4 m at the summit) during the time of 12:30-13:00 (Fig. 1b, Supplement Table S1).

60 The transmission velocity is the most critical parameter for estimating snowpack thickness. Because of the limited measurement time window in the so-called ‘death zone’, we did not measure common midpoint data to evaluate the transmission velocity of radar waves inside the snowpack at the Mount Everest. In general, the transmission velocity in snow ranges from 0.20 m/ns to 0.27 m/ns, which depends on the snow properties (Kovacs et al., 1995; Fortin and Fortier,

2001; Singh et al., 2017). A transmission velocity of 0.23 m/ns was obtained in a snowpack according to radar measurements with a steel stake (40 cm in length and 2 cm in diameter) that was buried in ~~the~~ snowpack at elevations of ~~6500 m and~~ 7028 m in 2005 (Sun et al., 2006). Therefore, we adopted a mean transmission velocity of 0.23 m/ns in this study.

To produce radar images that were more suitable for straightforward interpretations, the raw GPR data were processed using the Sensors & Software EKKO\_project processing package by ~~applying~~ ~~apply~~ a frequency bandpassing filter and time-variable gain corrections. The processing steps increased ~~the~~ signal-to-noise ratio to improve the imaging results while maintaining the original data signature, thus producing data that can be easily interpreted. The boundary between the snow and rock and the subsurface stratigraphies were visually traced.

### 3 Results and discussion

The radar wavelet traces showed strong signal contrast between the snow and the rock surface (the ~~blue red~~-dashed curve in Fig. 1c). It displays a well-defined gradual trend of radar reflection along the direction from the exposed limestone to ~~the Mount Everest~~ ~~the summit~~ (from wavelet No.1 to ~~wavelet~~ No.31), which indicates ~~the~~ thickening inclination of snow ~~depth~~ ~~thickness~~ along the northern slope of ~~the~~ Mount ~~Chomolungma~~ Everest. Such a thickening pattern agrees with the observed thick snowpack exposed by the nearby cliff and the topographic conditions for snow accumulation (Fig. 1a). ~~It should be noted that such a measurement along the north slope was used only for the purpose of generating the post-discerning radar boundaries, and the measurement process could give different results if the measurement profile were moved a few metres to either side.~~ The radar wavelet traces of the other 26 measuring points (Nos. 32-57), which ~~are~~ mainly concentrated at ~~the summit~~ ~~the Mount Everest~~ (Fig. 1b), displayed ~~the~~ similar radar reflections. Such homogeneity not only indicates the reliability of repeated radar measurements within this limited area, but also ~~provide~~ insights ~~into~~ the relatively flat topography along the ridge of ~~the~~ Mount Everest.

The magnitude of the estimated snow depth at ~~the~~ Mount Everest greatly depends on the choice of the mean transmission velocity. Taking the mean snow transmission velocity of 0.23 m/ns obtained at ~~6500 m asl and~~ 7028 m asl on ~~the~~ Mount ~~Chomolungma~~ Everest (Sun et al., 2006), we obtained the snow ~~depth~~ ~~thickness~~ distribution from the starting measurement point to the summit (Fig. 1c). The maximum two-way travel time of the reflecting horizon of the rock surface was approximately 88 ns at ~~the~~ Mount Everest. The snow ~~depth~~ ~~thickness~~ estimates gradually increased from ~2.0 m near the start of ~~the~~ exposed limestone to a maximum of ~10.1 m along the north slope. The snow ~~depth~~ ~~thickness~~ of a total of 26 measuring points concentrated at the summit was averaged to be approximately 9.5 m. Such thick accumulated snowpack at ~~the~~ Mount Everest may be partially explained by the westerly-introduced snowfall accumulation on the eastern leeward side. Moreover, compared with the ~~less~~ ~~lower~~ amount of snow accumulation ~~in~~ ~~along~~ the unfavourable steep slope, our radar ~~image~~ ~~measurements~~ covered the relatively flat platform at ~~the~~ Mount Everest (~~Fig. 1e~~), which may ~~provided~~ ~~the~~ favourable topography for snow accumulation.

Although the adopted transmission velocity in snow was determined at elevations of ~~6500 m and~~ 7028 m ~~in~~ Mount  
95 ~~Chomolungma Everest~~, ~~there may be still~~ some uncertainties ~~may still be~~ introduced by the distinct snow conditions ~~at the~~  
Mount Everest (e.g. ~~the~~ snow density ~~and~~; snow properties). The colder air temperature and stronger wind ~~levels~~ at higher  
elevations may ~~favor~~ ~~favour~~ ~~the~~ significant morphological changes, ~~by strong sublimation~~ and thus the snowpack was  
compacted, ~~resulting in~~ ~~for producing~~ high snow density. Therefore, if a higher ~~mean~~ snow density of ~~~5000.5~~  $\text{kg}/\text{m}^3$  were  
assumed ~~at the~~ Mount Everest, the mean transmission velocity would ~~reduced~~ ~~decrease~~ to ~~be~~  $\sim 0.21$  m/ns (Fortin and Fortier,  
100 2001). The mean snow ~~depth~~ ~~thickness~~ at ~~the~~ Mount Everest ~~will~~ ~~would~~ slightly ~~reduced~~ ~~decrease~~ from  $\sim 9.5$  m to  $\sim 8.7$  m. ~~The~~  
~~transmission velocity in snow generally ranges from 0.20 m/ns to 0.27 m/ns (Kovacs et al., 1995; Fortin and Fortier, 2001;~~  
~~Singh et al., 2017)~~ ~~The snow depth estimate will be changed by 0.4 m in 0.01 m/ns step of transmission velocity.~~ ~~Taking~~  
~~0.20 m/ns and 0.26 m/ns as the possible lower and upper boundaries for uncertainty estimation, the mean depth estimates at~~  
~~the summit were approximately  $9.5 \pm 1.2$  m in May 2022.~~

105 ~~In fact, the snow depth at Mount Everest should display interannual variability because of the influences of snow~~  
~~accumulation and snow drift. According to the recall of mountaineers who reached the summit in 2021 and 2022, the~~  
~~previously exposed rock surface in May 2021 was covered by snowpack of approximately 60-70 cm in May 2022. Our~~  
~~reported snow depth for Mount Everest in 2022 is considerably deeper than the values that were previously reported during~~  
~~the past five decades (0.9~3.5 m). There is still a lack of evidence that the snowpack has become thicker or thinner in recent~~  
110 ~~decades. Future repeated radar measurements at the summit would be helpful for evidencing such dynamic changes under~~  
~~climate change.~~

In addition to revealing the magnitude of ~~the~~ snow ~~depth~~ ~~thickness~~ at ~~the~~ Mount Everest ~~in May 2022~~, the radar wavelet  
traces showed two possible subsurface reflections within the snowpack (the yellow dashed lines in Fig. 1c). The upper weak  
subsurface reflection displays a shallow trend from a burial depth of  $\sim 2$ -3 m along the north slope to  $\sim 0.8$ -1.0 m. Another  
115 weak reflection layer ~~was~~ existed at a relatively uniform depth of ~~approximately~~ ~~appropriately~~ 4.5 m (Fig.1c). Such features  
may ~~be~~ attributed to the transition boundary ~~iesy~~ between fresh snow, compacted older snow and granular firn. However, this  
remains speculative due to the weak signal contrasts between layers.

#### 4 Conclusions

Overall, our measurements ~~acquired~~ in May 2022 provide the first clear radar image of ~~the~~ snowpack at the ~~summit~~ ~~top~~ of  
120 Mount Everest ~~in the world~~. This updated snow ~~depth~~ ~~thickness~~ at ~~the~~ Mount Everest is considerably deeper than  
~~the~~ ~~previously reported~~ values ~~that were previously reported~~ during the past five decades (0.9~3.5 m). ~~Such efforts~~  
~~gives~~ ~~provide the~~ ~~new insights for deciphering the true rock height and bottom geomorphology of Mount Everest. It is worth~~  
~~noting that recent debates took place on the surface melting that occurred at extremely high elevations (above 8000 m asl) on~~  
~~Mount~~ ~~in the Everest (Brun et al., 2022; Potocki et al., 2022).~~ ~~Such effort gives the new insights for not only deciphering the~~  
125 ~~true rock height and bottom geomorphology of the Mount Everest, but also for projecting future dynamic changes in the~~

~~cryosphere at such extremely high elevations due to anthropogenic warming.~~ It is worth noting that recent debates on the surface melting occurred at extremely high elevations (above 8000m asl) in the Everest (Brun et al., 2022; Potocki et al., 2022). Indeed, future snow core drilling and repeated ground penetrating radar measurements at ~~the~~ Mount Everest ~~are~~ also necessary to not only increase our understanding of ~~snow~~ dynamic ~~snow~~ changes, but also ~~to detect~~ ~~favor~~ ~~for~~ ~~detecting~~ the possible influence of ~~unprecedented~~ anthropogenic ~~climate changes~~ ~~source~~ ~~warming~~ by exploring the snow stratigraphy and ~~snowpack properties~~ ~~dynamic~~ ~~thickness~~ ~~changes~~ at the Earth's summit.

## References

- Angus-Leppan, P.: The height of mount Everest, *Surv. Rev.*, 26, 367-385, <https://doi.org/10.1179/sre.1982.26.206.367>, 1982.
- Brun, F., King, O., Réveillet, M., Amory, C., Planchot, A., Berthier, E., Dehecq, A., Bolch, T., Fourteau, K., and Brondex, J.: Brief communication: Everest South Col Glacier did not thin during the last three decades, *The Cryosphere Discussions*, 2022, 1-27, <https://doi.org/10.5194/tc-2022-166>, 2022.
- Chen, J., Zhang, Y., Yuan, J., Guo, C., and Zhang, P.: Height Determination of Qomolangma Feng (MT. Everest) in 2005, *Surv. Rev.*, 42, 122-131, <https://doi.org/10.1179/003962610X12572516251565>, 2010.
- Fortin, R. and Fortier, R.: Tomographic imaging of a snowpack, *Proceedings of the 58th Annual Eastern Snow Conference*, 2001, 17-19.
- Holbrook, W. S., Miller, S. N., and Provart, M. A.: Estimating snow water equivalent over long mountain transects using snowmobile-mounted ground-penetrating radar, *Geophys.*, 81, WA183-WA193, <https://doi.org/10.1190/geo2015-0121.1>, 2016.
- Hugonnet, R., McNabb, R., Berthier, E., Menounos, B., Nuth, C., Girod, L., Farinotti, D., Huss, M., Dussailant, I., and Brun, F.: Accelerated global glacier mass loss in the early twenty-first century, *Nature*, 592, 726-731, <https://doi.org/10.1038/s41586-021-03436-z>, 2021.
- Immerzeel, W. W., Lutz, A., Andrade, M., Bahl, A., Biemans, H., Bolch, T., Hyde, S., Brumby, S., Davies, B., and Elmore, A.: Importance and vulnerability of the world's water towers, *Nature*, 577, 364-369, <https://doi.org/10.1038/s41586-019-1822-y>, 2020.
- Kang, S., Zhang, Q., Zhang, Y., Guo, W., Ji, Z., Shen, M., Wang, S., Wang, X., Tripathee, L., and Liu, Y.: Warming and thawing in the Mt. Everest region: A review of climate and environmental changes, *Earth. Sci. Rev.*, 225, 103911, <https://doi.org/10.1016/j.earscirev.2021.103911>, 2022.
- Kraaijenbrink, P. D. A., Stigter, E. E., Yao, T., and Immerzeel, W. W.: Climate change decisive for Asia's snow meltwater supply, *Nat. Clim. Change.*, 11, 591-597, <https://doi.org/10.1038/s41558-021-01074-x>, 2021.
- Kovacs, A., Gow, A. J., and Morey, R. M.: The in-situ dielectric constant of polar firn revisited, *Cold. Reg. Sci. Technol.*, 23, 245-256, [https://doi.org/10.1016/0165-232X\(94\)00016-Q](https://doi.org/10.1016/0165-232X(94)00016-Q), 1995.
- Matthews, T., Perry, L. B., Koch, I., Aryal, D., Khadka, A., Shrestha, D., Abernathy, K., Elmore, A., Seimon, A., and Tait, A.: Going to extremes: installing the world's highest weather stations on Mount Everest, *Bull. Am. Meteorol. Soc.*, 101, E1870-E1890, <https://doi.org/10.1175/BAMS-D-19-0198.1>, 2020.
- Pepin, N., Adler, C., Kotlarski, S., and Palazzi, E.: Mountains undergo enhanced impacts of climate change, *Eos Earth & Space Science News*, 103, <https://doi.org/10.1029/2022EO185278>, 2022.
- Potocki, M., Mayewski, P. A., Matthews, T., Perry, L. B., Schwikowski, M., Tait, A. M., Korotkikh, E., Clifford, H., Kang, S., and Sherpa, T. C.: Mt. Everest's highest glacier is a sentinel for accelerating ice loss, *npj. Clim. Atmos. Sci.*, 5, 1-8, <https://doi.org/10.1038/s41612-022-00230-0>, 2022.

- 165 [Rignot, E., Mouginot, J., Larsen, C., Gim, Y., and Kirchner, D.: Low-frequency radar sounding of temperate ice masses in Southern Alaska, \*Geophys. Res. Lett.\*, 40, 5399-5405, <https://doi.org/10.1002/2013GL057452>, 2013.](#)
- [Singh, K., Negi, H., Kumar, A., Kulkarni, A., Dewali, S., Datt, P., Ganju, A., and Kumar, S.: Estimation of snow accumulation on Samudra Tapu glacier, Western Himalaya using airborne ground penetrating radar, \*Cur. Sci.\*, 2017. 1208-1218, <http://dx.doi.org/10.18520/cs/v112/i06/1208-1218>, 2017.](#)
- 170 [Sun, Z., Zhang, J.-q., and Zhang, P.: The application of radar technique to determining niveal bed thickness of mount Qomolangma, \*Geophysical and Geochemical Exploration\*, 2006. 179-182, 2006.](#)
- [Thompson, L. G., Brecher, H. H., Mosley-Thompson, E., Hardy, D. R., and Mark, B. G.: Glacier loss on Kilimanjaro continues unabated, \*Proc. Natl. Acad. Sci. U.S.A.\*, 106, 19770-19775, <http://dx.doi.org/10.1073/pnas.0906029106>, 2009.](#)
- [Xie, Y., Shen, W., Han, J., and Deng, X.: Determination of the height of Mount Everest using the shallow layer method, \*Geod. Geodyn.\*, 12, 258-265, <https://doi.org/10.1016/j.geog.2021.04.002>, 2021.](#)
- 175 [Yamamoto, T., Matsuoka, K., and Naruse, R.: Observation of internal structures of snow covers with a ground-penetrating radar, \*Ann. Glaciol.\*, 38, 21-24, <https://doi.org/10.3189/172756404781814834>, 2004.](#)
- [Angus-Leppan, P.: The height of mount everest, \*Survey Review\*, 26, 367-385, 1982.](#)
- [Brun, F., King, O., Réveillet, M., Amory, C., Planchot, A., Berthier, E., Dehecq, A., Bolch, T., Fourteau, K., and Brondex, J.: Brief communication: Everest South Col Glacier did not thin during the last three decades, \*The Cryosphere Discussions\*, 2022. 1-27, 2022.](#)
- 180 [Chen, J., Zhang, Y., Yuan, J., Guo, C., and Zhang, P.: Height Determination of Qomolangma Feng \(Mt. Everest\) in 2005, \*Survey Review\*, 42, 122-131, 2010.](#)
- [Fortin, R. and Fortier, R.: Tomographic imaging of a snowpack, 2001, 17-19.](#)
- 185 [GSSI, i.: GSSI Equipment at the Top of Mount Everest, <https://www.geophysical.com/gssi-equipment-at-the-top-of-mount-everest>, 2021. 2021.](#)
- [Holbrook, W. S., Miller, S. N., and Provart, M. A.: Estimating snow water equivalent over long mountain transects using snowmobile-mounted ground-penetrating radar, \*Geophysics\*, 81, WA183-WA193, 2016.](#)
- [Hugonnet, R., McNabb, R., Berthier, E., Menounos, B., Nuth, C., Girod, L., Farinotti, D., Huss, M., Dussaillant, L., and Brun, F.: Accelerated global glacier mass loss in the early twenty-first century, \*Nature\*, 592, 726-731, 2021.](#)
- 190 [Immerzeel, W. W., Lutz, A., Andrade, M., Bahl, A., Biemans, H., Bolch, T., Hyde, S., Brumby, S., Davies, B., and Elmore, A.: Importance and vulnerability of the world's water towers, \*Nature\*, 577, 364-369, 2020.](#)
- [Kang, S., Zhang, Q., Zhang, Y., Guo, W., Ji, Z., Shen, M., Wang, S., Wang, X., Tripathee, L., and Liu, Y.: Warming and thawing in the Mt. Everest region: A review of climate and environmental changes, \*Earth Science Reviews\*, 225, 103911, 2022.](#)
- 195 [Kraaijenbrink, P. D. A., Stigter, E. E., Yao, T., and Immerzeel, W. W.: Climate change decisive for Asia's snow meltwater supply, \*Nature Climate Change\*, 11, 591-597, 2021.](#)
- [Matthews, T., Perry, L. B., Koch, I., Aryal, D., Khadka, A., Shrestha, D., Abernathy, K., Elmore, A., Seimon, A., and Tait, A.: Going to extremes: installing the world's highest weather stations on Mount Everest, \*Bulletin of the American Meteorological Society\*, 101, E1870-E1890, 2020.](#)
- 200 [Pepin, N., Adler, C., Kotlarksi, S., and Palazzi, E.: Mountains undergo enhanced impacts of climate change, \*Eos Earth & Space Science News\*, 103, 2022.](#)
- [Potocki, M., Mayewski, P. A., Matthews, T., Perry, L. B., Schwikowski, M., Tait, A. M., Korotkikh, E., Clifford, H., Kang, S., and Sherpa, T. C.: Mt. Everest's highest glacier is a sentinel for accelerating ice loss, \*npj Climate and Atmospheric Science\*, 5, 1-8, 2022.](#)
- 205 [Rignot, E., Mouginot, J., Larsen, C., Gim, Y., and Kirchner, D.: Low-frequency radar sounding of temperate ice masses in Southern Alaska, \*Geophysical Research Letters\*, 40, 5399-5405, 2013.](#)
- [Singh, K., Negi, H., Kumar, A., Kulkarni, A., Dewali, S., Datt, P., Ganju, A., and Kumar, S.: Estimation of snow accumulation on Samudra Tapu glacier, Western Himalaya using airborne ground penetrating radar, \*Current Science\*, 2017. 1208-1218, 2017.](#)

- 210 ~~Sun, Z., Zhang, J. q., and Zhang, P.: The application of radar technique to determining niveal bed thickness of mount Qomolangma, Geophysical and Geochemical Exploration, 2006. 179-182, 2006.~~  
~~Thompson, L. G., Brecher, H. H., Mosley Thompson, E., Hardy, D. R., and Mark, B. G.: Glacier loss on Kilimanjaro continues unabated, Proceedings of the National Academy of Sciences, 106, 19770-19775, 2009.~~  
215 ~~Tone, S.: Measuring Mount Everest, the Rooftop of the World, <https://www.sixthtone.com/news/1005695/measuring-mount-everest%2C-the-rooftop-of-the-world>, 2020. 2020.~~  
~~Xie, Y., Shen, W., Han, J., and Deng, X.: Determination of the height of Mount Everest using the shallow layer method, Geodesy and Geodynamics, 12, 258-265, 2021.~~  
~~Yamamoto, T., Matsuoka, K., and Naruse, R.: Observation of internal structures of snow covers with a ground-penetrating radar, Annals of Glaciology, 38, 21-24, 2004.~~  
220

### **Data availability**

The GPR data (.sgy) are available from the corresponding author on reasonable request.

### **Author Contributions**

- 225 T.Y. designed research; T.Y., H.Z. and W.Y. analyzed data; and T.Y., H.Z., W.Y., B.X., J.Li., W.W., G.Wu and Z.W. jointly discuss the result and wrote the paper.

### **Acknowledgements**



The study is supported by the Second Tibetan Plateau Scientific Expedition and Research Program (STEP) (2019QZKK0201), the National Natural Science Foundation of China (41961134035).



### **230 Competing interests**

The authors declare no competing financial interests.









**Figure 1.** Radar measurements along the north slope to the Mount Everest acquired on May 4th, 2022. (a) Photo of Mount Everest showing the summit topography in 2022 and the radar measurement direction, as viewed from the northeast. (b) Distribution of 57 radar measurement points (red triangles), which started at the downwards-exposed metamorphosed limestone. (c) Radar wavelet traces showing the boundary between the snow and rock (red-blue dashed line) and the possible internal stratigraphies (yellow dashed lines) along the radar measurement profile at the estimated depth according to a constant transmission velocity (left axis) and the two-way wave travel time (right axis).